

## PRODUCT MONOGRAPH

Pr **PREMARIN**<sup>®</sup> Vaginal Cream  
(Conjugated Estrogens CSD, 0.625 mg/g)

### ESTROGENIC HORMONES

©Wyeth Canada.  
1025 Marcel Laurin Blvd.  
St. Laurent (Montreal), Quebec  
Canada H4R 1J6

Date of Revision:  
**June 01, 2007**

Submission Control No: 109701

## Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION .....	3
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS.....	3
WARNINGS AND PRECAUTIONS .....	5
ADVERSE REACTIONS .....	15
<b>DRUG INTERACTIONS</b> .....	20
DOSAGE AND ADMINISTRATION .....	23
OVERDOSAGE.....	24
ACTION AND CLINICAL PHARMACOLOGY .....	25
STORAGE AND STABILITY .....	28
SPECIAL HANDLING INSTRUCTIONS.....	28
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	29
PART II: SCIENTIFIC INFORMATION.....	30
PHARMACEUTICAL INFORMATION .....	30
CLINICAL TRIALS .....	30
DETAILED PHARMACOLOGY .....	31
REFERENCES .....	33
PART III: CONSUMER INFORMATION.....	37

**Premarin® Vaginal Cream**  
(Conjugated Estrogens CSD, 0.625 mg/g)

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
vaginal	conjugated estrogens vaginal cream 0.625 mg/g	<i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

**INDICATIONS AND CLINICAL USE**

**Premarin® Vaginal Cream is indicated in the treatment of atrophic vaginitis, dyspareunia, and kraurosis vulvae.**

**PREMARIN® VAGINAL CREAM HAS NOT BEEN SHOWN TO BE EFFECTIVE FOR ANY PURPOSE DURING PREGNANCY AND ITS USE MAY CAUSE SEVERE HARM TO THE FETUS.**

**Premarin® Vaginal Cream should be prescribed with an appropriate dose of a progestin for women with intact uteri in order to prevent endometrial hyperplasia/carcinoma**

**Geriatrics (> 65 years of age):** See above indications.

**Pediatrics (< 16 years of age):** Premarin® Vaginal Cream is not indicated for use in children.

**CONTRAINDICATIONS**

Premarin® Vaginal Cream should not be administered to patients with any of the following conditions:

- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.
- Liver dysfunction or disease as long as liver function tests have failed to return to normal.
- Known or suspected estrogen-dependent malignant neoplasia (e.g. endometrial cancer).
- Endometrial hyperplasia.
- Known, suspected, or past history of breast cancer.
- Undiagnosed abnormal genital bleeding.
- Known or suspected pregnancy.
- Active or past history of confirmed venous thromboembolism (such as deep venous thrombosis or pulmonary embolism) or active thrombophlebitis.
- Active or past history of arterial thromboembolic disease (e.g. stroke, myocardial infarction, coronary heart disease).
- Partial or complete loss of vision due to ophthalmic vascular disease.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

The Women's Health Initiative (WHI) trial examined the health benefits and risks of oral combined *estrogen plus progestin* therapy (n=16,608) and oral *estrogen-alone* therapy (n=10,739) in postmenopausal women aged 50 to 79 years.<sup>3, 6, 42</sup>

The *estrogen plus progestin* arm of the WHI trial (mean age 63.3 years) indicated an increased risk of *myocardial infarction* (MI), *stroke*, *invasive breast cancer*, *pulmonary emboli* and *deep vein thrombosis* in postmenopausal women receiving treatment with combined conjugated equine estrogens (CEE, 0.625 mg/day) and medroxyprogesterone acetate (MPA, 2.5 mg/day) for 5.2 years compared to those receiving placebo.<sup>42</sup>

The *estrogen-alone* arm of the WHI trial (mean age 63.6 years) indicated an increased risk of *stroke* and *deep vein thrombosis* in hysterectomized women treated with CEE-alone (0.625 mg/day) for 6.8 years compared to those receiving placebo.<sup>3</sup>

Therefore, the following should be given serious consideration at the time of prescribing:

- Estrogens with or without progestins **should not** be prescribed for primary or secondary prevention of cardiovascular diseases.
- Estrogens with or without progestins should be prescribed at **the lowest effective dose** for the approved indication.
- Estrogens with or without progestins should be prescribed for the **shortest period possible** for the approved indication.

### General

**Estrogen Therapy (ET) or Hormone Therapy (HT) should not be used for the prevention of cardiovascular disease or dementia.**

**Estrogen Replacement Therapy (ERT) and Hormone Replacement Therapy (HRT) have been associated with increased risks of certain cancers and cardiovascular diseases. The use of unopposed estrogens in women with intact uteri is associated with an increased risk of endometrial cancer.**

**The benefits and risks of ERT and HRT must always be carefully weighed, including**

**consideration of the emergence of risks as therapy continues. Estrogens with or without progestins should be prescribed at the lowest effective doses and for the shortest duration consistent with treatment goals and risks for the individual woman.** In the absence of comparable data, the risks of HRT should be assumed to be similar for all estrogens and estrogen/progestin combinations.

Systematic absorption may occur with the use of conjugated estrogens (CE) vaginal cream. Warnings, precautions and adverse reactions associated with oral CE treatment should be taken into account.

### **Latex Condoms**

**NOTE:** Preliminary studies conducted by the Health Products and Food Branch have demonstrated that PREMARIN® Vaginal Cream may react with the latex rubber of certain mechanical barrier devices used for prevention of sexually transmitted diseases and pregnancy (diaphragms and condoms). In additional studies, Premarin Vaginal Cream has been shown to weaken latex condoms. The potential for Premarin Vaginal Cream to weaken and contribute to the failure of condoms, diaphragms, or cervical caps made of latex or rubber should be considered.

### **Carcinogenesis and Mutagenesis**

#### ***Breast cancer***

Available epidemiological data indicate that the use of combined *estrogen plus progestin* by postmenopausal women is associated with an increased risk of invasive breast cancer.

In the *estrogen plus progestin* arm of the WHI trial, among 10,000 women over a one-year period, there were:

- 8 more cases of invasive breast cancer (38 on combined HRT versus 30 on placebo).<sup>42</sup>

The WHI study also reported that the invasive breast cancers diagnosed in the *estrogen plus progestin* group were similar in histology but were larger (mean [SD], 1.7 cm [1.1] vs 1.5 cm [0.9], respectively; P=0.04) and were at a more advanced stage compared with those diagnosed in the placebo group. The percentage of women with abnormal mammograms (recommendations for short-interval follow-up, a suspicious abnormality, or highly suggestive of malignancy) was significantly higher in the *estrogen plus progestin* group versus the placebo group. This difference appeared at year one and persisted in each year thereafter.<sup>6</sup>

In the *estrogen-alone* arm of the WHI trial, there was no statistically significant difference in the rate of invasive breast cancer in hysterectomized women treated with conjugated equine estrogens versus women treated with placebo.<sup>3</sup>

It is recommended that estrogens not be given to women with existing breast cancer or those with a previous history of the disease (see **CONTRAINDICATIONS**).

There is a need for caution in prescribing estrogens for women with known risk factors associated with the development of breast cancer, such as strong family history of breast cancer (first degree relative) or who present a breast condition with an increased risk (abnormal mammograms and/or atypical hyperplasia at breast biopsy).

Other known risk factors for the development of breast cancer such as nulliparity, obesity, early menarche, late age at first full term pregnancy and at menopause should also be evaluated.

It is recommended that women undergo mammography prior to the start of HRT treatment and at regular intervals during treatment, as deemed appropriate by the treating physician and according to the perceived risks for each patient.

The overall benefits and possible risks of hormone replacement therapy should be fully considered and discussed with patients. It is important that the modest increased risk of being diagnosed with breast cancer after 4 years of treatment with combined estrogen plus progestin HRT (as reported in the results of the WHI trial) be discussed with the patient and weighed against its known benefits.

*Instructions for regular self-examination of the breasts should be included in this counselling.*

### ***Endometrial hyperplasia & endometrial carcinoma***

Estrogen-only HRT increases the risk of endometrial hyperplasia/carcinoma if taken by women with intact uteri. Estrogen should be prescribed with an appropriate dosage of a progestin for women with intact uteri in order to prevent endometrial hyperplasia/carcinoma.

The reported endometrial cancer risk among unopposed estrogen users is about 2- to 12-fold or greater than in non-users and appears to be dependent on duration of treatment and on estrogen dose. Most studies show no significant increased risk associated with the use of estrogens for less than one year. The greatest risk appears associated with prolonged use, with increased risks of 15- to 24-fold for five years or more, and this risk has been shown to persist for at least 8 to 15 years after ERT is discontinued.

In a subset of WHI (see ACTIONS AND CLINICAL PHARMACOLOGY) no increased risk of endometrial cancer after an average of 5.2 years of treatment with combined estrogen plus progestin HRT compared to placebo was observed.

Endometrial hyperplasia (a possible precursor of endometrial cancer) has been reported to occur at a rate of approximately 1% or less with CEE or CEE/MPA in two large clinical trials [Health and Osteoporosis, Progestin and Estrogen (n=2,153) and Menopausal Study Group (n=1,385)].<sup>28, 40</sup> In these two clinical trials two cases of endometrial cancer were reported to occur among women taking combination CEE/MPA.

Clinical surveillance of all women taking combined estrogen plus progestin HRT is important. Adequate diagnostic measures, including endometrial sampling when indicated, should be undertaken to rule out malignancy in all cases of undiagnosed persistent or recurring abnormal vaginal bleeding.

### ***Ovarian cancer***

In some epidemiologic studies, use of estrogen-only products, in particular for ten or more years, has been associated with an increased risk of ovarian cancer. The analysis of the WHI data suggested that estrogen plus progestin therapy may increase the risk of ovarian cancer. Other epidemiologic studies have not found these associations.<sup>2, 18, 31</sup>

## **Cardiovascular**

### ***Cardiovascular risk***

Oral HRT has been associated with an increased risk of cardiovascular events such as myocardial infarction (MI) and stroke, as well as venous thrombosis and pulmonary embolism (PE) (venous thromboembolism or VTE). Should any of these occur or be suspected, HRT should be discontinued immediately.

Risk factors for cardiovascular disease (e.g., hypertension, diabetes mellitus, tobacco use, hypercholesterolemia, and obesity) should be managed appropriately.

### ***General***

The results of the Heart and Estrogen/progestin Replacement Studies (HERS and HERS II) and the Women's Health Initiative (WHI) trial indicate that the use of *estrogen plus progestin* is associated with an increased risk of coronary heart disease (CHD) in postmenopausal women.<sup>9, 16, 42</sup> The results of the WHI trial indicate that the use of *estrogen-alone* and *estrogen plus progestin* is associated with an increased risk of stroke in postmenopausal women.<sup>3, 42</sup>

### ***WHI trial findings***

In the combined *estrogen plus progestin* arm of the WHI trial, among 10,000 women over a one-year period, there were:

- 8 more cases of stroke (29 on combined HRT versus 21 on placebo).
- 7 more cases of CHD (37 on combined HRT versus 30 on placebo).<sup>42</sup>

In the *estrogen-alone* arm of the WHI trial of women with prior hysterectomy, among 10,000 women over a one-year period, there were/was:

- 12 more cases of stroke (44 on *estrogen-alone* therapy versus 32 on placebo)
- no statistically significant difference in the rate of CHD.<sup>3</sup>

### ***HERS and HERS II findings***

In the Heart and Estrogen/progestin Replacement Study (HERS) of postmenopausal women with documented heart disease (n=2763, average age 66.7 years), a randomized placebo-controlled clinical trial of secondary prevention of coronary heart disease (CHD), treatment with 0.625 mg/day oral conjugated equine estrogen (CEE) plus 2.5 mg oral medroxyprogesterone acetate (MPA) demonstrated no cardiovascular benefit. Specifically, during an average follow-up of 4.1 years, treatment with CEE plus MPA did not reduce the overall rate of CHD events in postmenopausal women with established coronary heart disease. There were more CHD events in the hormone-treated group than in the placebo group in year 1, but not during the subsequent years.<sup>16</sup>

From the original HERS trial, 2321 women consented to participate in an open label extension of HERS known as HERS II. Average follow-up in HERS II was an additional 2.7 years, for a total of 6.8 years overall. After 6.8 years, hormone therapy did not reduce the risk of cardiovascular events in women with CHD.<sup>9</sup>

### ***Blood pressure***

Women using hormone replacement therapy sometimes experience increased blood pressure. Blood pressure should be monitored with HRT use. Elevation of blood pressure in previously normotensive or hypertensive patients should be investigated and HRT may have to be discontinued.

### **Endocrine and Metabolism**

#### ***Glucose and lipid metabolism***

A worsening of glucose tolerance and lipid metabolism have been observed in a significant percentage of peri- and post-menopausal patients. Therefore, diabetic patients, or those with a

predisposition to diabetes, should be observed closely to detect any alterations in carbohydrate or lipid metabolism, especially in triglyceride blood levels.

Women with familial hyperlipidemias need special surveillance. Lipid-lowering measures are recommended additionally, before treatment is started.

Caution should be exercised in patients with pre-existing hypertriglyceridemia since rare cases of large increases of plasma triglycerides leading to pancreatitis have been reported with estrogen therapy in this population.

Women with porphyria need special surveillance.

Estrogens should be used with caution in individuals with severe hypocalcemia.

### **Calcium and phosphorus metabolism**

Because the prolonged use of estrogens influences the metabolism of calcium and phosphorus, estrogens should be used with caution in patients with metabolic and malignant bone diseases associated with hypercalcemia and in patients with renal insufficiency.

#### *Hypothyroidism*

Patients who require thyroid hormone replacement therapy and who are also taking estrogen should have their thyroid function monitored regularly to assure that thyroid hormone levels remain in an acceptable range (see *Drug-Laboratory Test Interactions*).

### **Genitourinary**

#### *Endometriosis*

Symptoms and physical findings associated with a previous diagnosis of endometriosis may reappear or become aggravated with estrogen use.

Addition of a progestin should be considered in women who have undergone a hysterectomy but are known to have residual endometriosis, since a few cases of malignant transformation after estrogen-only therapy have been reported.

#### *Uterine Leiomyomata*

Pre-existing uterine leiomyomata may increase in size during estrogen use. Growth, pain or tenderness of uterine leiomyomata requires discontinuation of medication and appropriate investigation.

### ***Vaginal bleeding***

Abnormal vaginal bleeding, due to its prolongation, irregularity or heaviness, occurring during therapy should prompt appropriate diagnostic measures to rule out the possibility of uterine malignancy and the treatment should be re-evaluated.

### **Hematologic**

#### *Venous thromboembolism*

Available epidemiological data indicate that use of estrogen with or without progestin by postmenopausal women is associated with an increased risk of developing venous thromboembolism (VTE).

In the oral *estrogen plus progestin* arm of the WHI trial, among 10,000 women on combined HRT over a one-year period, there were 18 more cases of venous thromboembolism, including 8 more cases of pulmonary embolism.<sup>42</sup>

In the oral *estrogen-alone* arm of the WHI trial, among 10,000 women on estrogen therapy over a one-year period, there were 7 more cases of venous thromboembolism, although there was no statistically significant difference in the rate of pulmonary embolism.<sup>3</sup>

Generally recognized risk factors for VTE include a personal history, a family history (the occurrence of VTE in a direct relative at a relatively early age may indicate genetic predisposition), severe obesity (body mass index > 30 kg/m<sup>2</sup>) and systemic lupus erythematosus. The risk of VTE also increases with age and smoking.

The risk of VTE may be temporarily increased with prolonged immobilization, major surgery or trauma. In women on HRT, attention should be given to prophylactic measures to prevent VTE following surgery. Also, patients with varicose veins should be closely supervised. The physician should be alert to the earliest manifestations of thrombotic disorders (thrombophlebitis, retinal thrombosis, cerebral embolism and pulmonary embolism). If these occur or are suspected, hormone therapy should be discontinued immediately, given the risks of long-term disability or fatality.

If feasible, estrogens should be discontinued at least 4 weeks before major surgery which may be associated with an increased risk of thromboembolism, or during periods of prolonged immobilization.

## **Hepatic/Biliary/Pancreatic**

### ***Gallbladder diseases***

A 2 to 4-fold increase in the risk of gallbladder disease requiring surgery in women receiving postmenopausal estrogens has been reported.

### ***Hepatic hemangiomas***

Particular caution is indicated in women with hepatic hemangiomas, as HRT may cause an exacerbation of this condition.

### ***Jaundice***

Caution is advised in patients with a history of liver and/or biliary disorders. If cholestatic jaundice develops during treatment, the treatment should be discontinued and appropriate investigations carried out.

### ***Liver function tests***

Liver function tests should be done periodically in subjects who are suspected of having hepatic disease. For information on endocrine and liver function tests, see **Monitoring and Laboratory Tests**.

## **Immune**

Particular caution is indicated in women with systemic lupus erythematosus, as HRT may cause an exacerbation of this condition.

## **Neurologic**

### ***Cerebrovascular insufficiency***

Patients who develop visual disturbances, classical migraine, transient aphasia, paralysis or loss of consciousness should discontinue medication.

Patients with a previous history of classical migraine and who develop a recurrence or worsening of migraine symptoms should be reevaluated.

### ***Dementia***

Available epidemiological data indicate that the use of combined *estrogen plus progestin* in women age 65 and over may increase the risk of developing probable dementia.

The Women's Health Initiative Memory Study (WHIMS), a clinical substudy of the WHI, was designed to assess whether postmenopausal hormone replacement therapy (oral *estrogen plus progestin* or oral *estrogen-alone*) reduces the risk of dementia in women aged 65 and over (age range 65-79 years) and free of dementia at baseline.<sup>32, 33</sup>

In the *estrogen plus progestin* arm of the WHIMS (n=4532), women with intact uteri were treated with daily 0.625 mg conjugated equine estrogens (CEE) plus 2.5 mg medroxyprogesterone acetate (MPA) or placebo for an average of 4.05 years. The results, when extrapolated to 10,000 women treated over a one-year period showed:

- 23 more cases of probable dementia (45 on combined HRT versus 22 on placebo).<sup>33</sup>

In the *estrogen-alone* arm of the WHIMS (n=2947), women with prior hysterectomy were treated with daily 0.625 mg CEE or placebo for an average of 5.21 years. The results, when extrapolated to 10,000 women treated over a one-year period showed:

- 12 more cases of probable dementia (37 on *estrogen-alone* versus 25 on placebo), although this difference did not reach statistical significance.<sup>32</sup>

When data from the *estrogen plus progestin* arm of the WHIMS and the *estrogen-alone* arm of the WHIMS were combined, as per the original WHIMS protocol, in 10,000 women over a one-year period, there were:

- 18 more cases of probable dementia (41 on *estrogen plus progestin* or *estrogen-alone* versus 23 on placebo).<sup>32</sup>

## **Epilepsy**

Particular caution is indicated in women with epilepsy, as HRT may cause an exacerbation of this condition.

## **Ophthalmologic**

### *Visual abnormalities*

Retinal vascular thrombosis has been reported in patients receiving estrogens. Discontinue medication pending examination if there is sudden partial or complete loss of vision, or a sudden onset of proptosis, diplopia, or migraine. If examination reveals papilledema or retinal vascular lesions, medication should be withdrawn.

## **Psychiatric**

### *Depression*

Patients who are taking progestogens and have a history of depression should be observed. If the depression occurs to a serious degree, the drug should be discontinued.

## **Renal**

### ***Fluid retention***

Estrogens may cause fluid retention.

Therefore, particular caution is indicated in cardiac, renal dysfunction, or asthma. If, in any of the above-mentioned conditions, a worsening of the underlying disease is diagnosed or suspected during treatment, the benefits and risks of treatment should be reassessed based on the individual case.

## **Special Populations**

**Pregnant Women:** Estrogens/progestins should not be used during pregnancy.

**Nursing Women:** Estrogen should not be used during lactation.

**Pediatrics (< 16 years of age):** Premarin<sup>®</sup> Vaginal Cream is not indicated for use in children.

**Geriatrics (> 65 years of age):** Of the total number of subjects in the conjugated equine estrogens in combination with medroxyprogesterone acetate substudy of the Women's Health Initiative study (WHI), 44% (n=7320) were 65 years and over, while 6.6% (n=1,095) were 75 and over. No significant differences in relative risks were observed between subjects 65 years and over compared to younger subjects. There was a higher relative risk of stroke and invasive breast cancer in women 75 and over compared to younger subjects.

## **Monitoring and Laboratory Tests**

Before Premarin<sup>®</sup> Vaginal Cream is administered, the patient should have a complete physical examination including blood pressure determination. Breasts and pelvic organs should be appropriately examined and a Papanicolaou smear should be performed. Endometrial biopsy should be done only when indicated. Baseline tests should include mammography, measurements of blood glucose, calcium, triglycerides and cholesterol, and liver function tests. Before starting treatment pregnancy should be excluded. Periodic check-ups and careful benefit/risk evaluations should be undertaken in women treated with ERT/HRT therapy. The

first follow-up examination should be done within three to six months of initiation of treatment to assess response to treatment. Thereafter, examinations should be made at intervals of at least once a year. Appropriate investigations should be arranged at regular intervals as determined by the physician.

Mammography examinations should be scheduled based on patient age, risk factors and prior mammogram results.

*The importance of regular self-examination of the breasts should be discussed with the patient.*

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

**See Warnings/Precautions regarding potential induction of malignant neoplasia and other adverse effects similar to those observed with oral contraceptives.**

**The following additional adverse reactions have been reported with conjugated estrogens vaginal cream or are undesirable effects associated with ET/HT:**

#### **Blood and lymphatic system disorders**

Altered coagulation tests (see **WARNINGS AND PRECAUTIONS**, *Drug-Laboratory Tests Interactions*).

#### **Cardiac disorders**

Palpitations; increase in blood pressure (see **WARNINGS AND PRECAUTIONS**); coronary thrombosis, pulmonary embolism, venous thrombosis, myocardial infarction.

#### **Endocrine disorders**

Increased blood sugar levels; decreased glucose tolerance, precocious puberty.

#### **Eye disorders**

Neuro-ocular lesions (e.g. retinal thrombosis, optic neuritis); visual disturbances; steepening of the corneal curvature; intolerance to contact lenses.

### **Gastrointestinal disorders**

Nausea; vomiting; abdominal discomfort (cramps, pressure, pain, bloating), pancreatitis.

### **General disorders and administration site conditions**

Fatigue; changes in appetite; changes in body weight; changes in libido, aggravation of porphyria, angioedema, hypersensitivity, anaphylactic/anaphylactoid reactions, increased triglycerides..

### **Hepatobiliary disorders**

Gallbladder disorder; cholestatic jaundice.

### **Musculoskeletal and connective tissue disorders**

Musculoskeletal pain including leg pain not related to thromboembolic disease (usually transient, lasting 3-6 weeks) may occur, arthralgias, leg cramps..

### **Nervous system disorders**

Aggravation of migraine episodes; headaches; migraines, dizziness; cerebrovascular accident/stroke.

### **Psychiatric disorders**

Mental depression; nervousness; irritability, mood disturbances, dementia.

### **Renal and urinary disorders**

Cystitis; dysuria; sodium retention; edema.

### **Reproductive system and breast disorders**

Breakthrough bleeding; spotting; change in menstrual flow; dysmenorrhea; vaginal itching/discharge; dyspareunia; endometrial hyperplasia; pre-menstrual-like syndrome; reactivation of endometriosis; changes in cervical erosion and amount of cervical secretion; breast swelling and tenderness, galactorrhea, breast discharge, amenorrhea, increase in size of uterine leiomyomata, vaginitis, application site reactions of vulvovaginal discomfort including burning, irritation and genital pruritus, vaginal candidiasis.

### **Skin and subcutaneous tissue disorders**

Chloasma or melasma, which may persist when drug is discontinued; erythema multiforme; erythema nodosum; haemorrhagic eruption; loss of scalp hair; hirsutism and acne, allergic reactions and rashes, generalized rash, urticaria, pigmentation of the skin, pruritis.

### **Vascular disorders**

Isolated cases of: thrombophlebitis; thromboembolic disorders.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse drug reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

A phase III double-blind, randomized study was conducted to compare the efficacy and safety of various regimens of Premarin® (conjugated estrogens) and medroxyprogesterone acetate (MPA). Efficacy was determined by the incidence of endometrial hyperplasia at the twelve month evaluation. A total of 1,724 generally healthy postmenopausal women (mean age, 54.0 years  $\pm$  SD 4.6) participated in the study. The patients were considered as having completed the study if they participated in all 13 cycles (28 days/cycle). The five arms in the study were: 2 for Premplus®, 2 for Premplus Cycle®, and 1 for Premarin® tablets alone.

Prior to treatment, the following were performed: physical examinations, vital signs, papanicolaou smear, laboratory safety screen, mammography, follicle stimulating hormone (FSH), and endometrial biopsy. During the patient visit for Cycle 6, all but the mammography and FSH were performed. At the end of the study, Cycle 13, all but the FSH were performed.

No dose-dependent incidence of adverse experiences was seen in the multicenter efficacy and safety study. Significantly ( $p < 0.05$ ) fewer (12%) Premarin® treated patients reported breast pain than in the Premarin®/MPA groups. Headache was the most common drug-related study event in the Premarin® alone group, reported by 69 (20%) patients. Table 1 summarizes the treatment-emergent drug-related study events reported by 2% or more of the patients.

Table 1: Treatment-Emergent Drug-Related Study Events With an Incidence of  $\geq 2\%$  for Premarin® Tablets.

Study Event	Premarin® 0.625 mg CE (n=347)  No. (%) of Patients <sup>+</sup>
<b>General disorders and administration site conditions</b>	
Asthenia	18 (5)
Chest pain	2 (<1)
Generalized edema	9 (3)
Edema	5 (1)
Peripheral edema	11 (3)
Pain	11 (3)
<b>Vascular disorders</b>	
Hypertension	7 (2)
Vasodilatation	9 (3)
<b>Gastrointestinal disorders</b>	
Diarrhea	6 (2)
Dyspepsia	4 (1)
Flatulence	14 (4) <sup>b</sup>
Nausea	19 (5)
Abdominal pain	46 (13)
<b>Musculoskeletal connective tissue, and bone disorders</b>	
Leg cramps	8 (2)
Back pain	13 (4)
<b>Nervous system disorders</b>	
Headache	69 (20)
Depression	22 (6)

Study Event	Premarin® 0.625 mg CE (n=347)  No. (%) of Patients <sup>+</sup>
Migraine	7 (2)
Dizziness	10 (3)
Emotional lability	4 (1)
Insomnia	2 (<1)
Nervousness	1 (<1) <sup>b, d</sup>
<b>Skin and subcutaneous tissue disorders</b>	
Acne	6 (2)
Pruritus	6 (2) <sup>a, b</sup>
Rash	5 (1)
<b>Reproductive system and breast disorders</b>	
Breast enlargement	4 (1) <sup>a, b</sup>
Breast pain *	40 (12) <sup>a, b, d</sup>
Cervix disorder**	12 (3)
Dysmenorrhea	17 (5) <sup>d</sup>
Endometrial hyperplasia	57 (20)
Leukorrhea	24 (7)
Menstrual disorder	3 (<1)
Pelvic pain	16 (5)
Uterine spasm	0 (0) <sup>a, d</sup>
Vaginal bleeding ***	28 (8) <sup>b</sup>
Vaginitis	4 (1) <sup>a, b</sup>
<b>Investigations</b>	
Pap smear abnormal <sup>†</sup>	0 (0) <sup>a</sup>

Study Event	Premarin® 0.625 mg CE (n=347)  No. (%) of Patients <sup>+</sup>
Weight increased	10 (3)
<b>Psychiatric disorders</b>	
Depression	22 (6)
Emotional lability	4 (1)
Nervousness	1 (<1) <sup>b,d</sup>

- + Patients were counted only once for a particular study event.
- \* Breast pain also includes breast discomfort, breast soreness, breast tenderness, mastodynia, nipple soreness and nipple tenderness.
- \*\* Cervix disorder includes cervical dysplasia, cervical erosion, cervical hypersecretion.
- † Pap smear abnormal refers to positive Pap smear class III through V.
- \*\*\* Vaginal bleeding includes menorrhagia, metrorrhagia, uterine hemorrhage, and vaginal hemorrhage.
- a, b, d, = Significant difference (  $p < 0.05$ ) from treatment group Premplus® (0.625/2.5 mg), Premplus® (0.625/5.0 mg), Premplus Cycle® (0.625/10.0 mg) and Premarin® (0.625 mg) respectively.

The above table summarizes the treatment-emergent drug-related study events reported by greater than 2% of the patients. The number of patients with any study event is not necessarily the sum of the individual events since a patient might have reported two or more different study events. The addition of progestin to estrogen replacement therapy may contribute to breast pain. This is reflected by the greater percentage of patients with breast pain on combination therapy than on Premarin alone.

If adverse symptoms persist, the prescription of HRT should be re-considered.

## **DRUG INTERACTIONS**

### **Overview**

In vitro and in vivo studies have shown that 17  $\beta$ -estradiol, one of the components of conjugated estrogens, is metabolized partially by Cytochrome P450 3A4 (CYP3A4). Therefore, strong CYP3A4 inducers such as phenobarbital, phenytoin, carbamazepine, rifampicin and dexamethasone may reduce plasma concentrations of 17  $\beta$ -estradiol. This may lead to a decreased effect and/or changes in the uterine bleeding profile. CYP3A4 inhibitors such as cimetidine, erythromycin, ketoconazole, clarithromycin, itraconazole, and ritonavir may increase plasma concentrations of 17  $\beta$ -estradiol and may result in side effects.

Data from a drug-drug interaction study involving conjugated estrogens and medroxyprogesterone acetate indicate that the pharmacokinetic disposition of both drugs are not altered when the drugs are co-administered. Other clinical drug-drug interaction studies have not been conducted with conjugated estrogens.

Estrogens may diminish the effectiveness of anticoagulant, antidiabetic and antihypertensive agents.

Preparations inducing liver enzymes (e.g. barbiturates, hydantoins, carbamazepine, meprobamates, phenylbutazone or rifampicin) may interfere with the activity of orally administered estrogens.

### **Drug-Drug Interactions**

The following section contains information on drug interactions with ethinyl estradiol-containing products (specifically, oral contraceptives) that have been reported in the public literature. **It is unknown whether such interactions occur with drug products containing other types of estrogens.**

**Hepatic metabolism:** Interactions can occur with drugs that induce microsomal enzymes which can decrease ethinyl estradiol concentrations (eg., rifampin, barbiturates, phenytoin, carbamazepine, troglitazone).

**Gastrointestinal wall:** Sulfation of ethinyl estradiol has been shown to occur in the gastrointestinal (GI) wall. Therefore, drugs which act as competitive inhibitors for sulfation in the GI wall may increase ethinyl estradiol bioavailability (eg., ascorbic acid, acetaminophen).

**Interference in the metabolism of other drugs:** Ethinyl estradiol may interfere with the metabolism of other drugs by inhibiting hepatic microsomal enzymes or by inducing hepatic drug conjugation, particularly glucuronidation. Increased plasma concentrations of cyclosporin, prednisolone, and theophylline have been reported with concomitant administration of certain drugs containing ethinyl estradiol (eg., oral contraceptives containing ethinyl estradiol). In addition, products containing ethinyl estradiol may induce the conjugation of other compounds.

Decreased plasma concentrations of acetaminophen and increased clearance of temazepam, salicylic acid, morphine, and clofibric acid have been noted when these drugs were administered with certain ethinyl estradiol-containing drug products (eg., oral contraceptives containing ethinyl estradiol).

**Other interactions with ethinyl estradiol:** Coadministration of atorvastatin and certain ethinyl estradiol-containing drug products (eg., oral contraceptives) increase AUC values for ethinyl estradiol by 20 percent.

Clinical pharmacokinetic studies have not demonstrated any consistent effect of antibiotics (other than rifampin) on plasma concentrations of synthetic steroids.

### **Drug-Food Interactions**

CYP3A4 inhibitors such as grapefruit juice may increase plasma concentrations of 17  $\beta$ -estradiol and may result in side effects.

### **Drug-Herb Interactions**

It was found that some herbal products (e.g., St. John's wort), which are available as over-the-counter (OTC) products, might interfere with steroid metabolism, and therefore alter the efficacy and safety of estrogen/progestin products. Hot flashes and vaginal bleeding have been reported in patients taking estrogen replacement therapy (ERT) and combined estrogen plus progestin therapy (HRT) and St. John's wort. St. John's wort may induce hepatic microsomal enzymes, which theoretically may result in reduced efficacy of ERT and HRT.

Physicians and other health care providers should be made aware of other non-prescription products concomitantly used by the patient, including herbal and natural products, obtained from the widely spread Health Stores.

### **Drug-Laboratory Test Interactions**

The results of certain endocrine and liver function tests may be affected by estrogen-containing products:

- increased prothrombin time and partial thromboplastin time; increased levels of fibrinogen and fibrinogen activity, increased coagulation factors VII, VIII, IX, X; increased norepinephrine-induced platelet aggregability; decreased antithrombin III;
- impaired glucose tolerance;
- increased serum triglycerides and phospholipids concentration.
- increased thyroid-binding globulin (TBG) levels leading to increased circulating total thyroid hormone ( $T_4$ ), as measured by column or radioimmunoassay;  $T_3$  resin uptake is decreased, reflecting the elevated TBG; free  $T_4$  concentration is unaltered.
- other binding proteins may be elevated in serum i.e., corticosteroid binding globulin (CBG), sex-hormone binding globulin (SHBG), leading to increased circulating corticosteroids and sex steroids respectively, free or biologically active hormone concentrations are unchanged;

The results of the above laboratory tests should not be considered reliable unless therapy has been discontinued for two to four weeks.

The pathologist should be informed that the patient is receiving hormone replacement therapy **(HRT) when relevant specimens are submitted.**

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Use of Premarin® Vaginal Cream, alone, or in combination with a progestin, should be limited to the shortest duration consistent with treatment goals and risks for the individual woman. Patients should be re-evaluated periodically to determine if treatment for symptoms is still necessary.

### **Recommended Dose and Dosage Adjustment**

Premarin® Vaginal Cream should be administered cyclically for short-term use only for the treatment of atrophic vaginitis, dyspareunia or kraurosis vulvae.

In patients with severe cases of atrophic vaginitis, the mucosa should first be conditioned with a short course of oral therapy -- 1.25 mg daily for approximately 10 days. Vaginal treatment should be instituted at the lowest effective dosage, and the requirement for estrogen therapy reassessed regularly.

In patients already receiving oral therapy, the oral dosage may be reduced taking into account the potential absorption from the vaginal medication. The degree of atrophy is directly responsible for the level of absorption and should be the guiding factor in dose adjustment.

In women with an intact uterus, a progestin should be coadministered for a minimum of 10, but preferably at least 12 to 14 days per cycle to avoid overstimulation of the endometrium. In addition, progestin should be administered to minimize the occurrence of endometrial hyperplasia. Unexpected or abnormal vaginal bleeding in such patients requires institution of prompt diagnostic measures, such as endometrial biopsy or curettage to rule out the possibility of uterine malignancy. Since progestins are administered to reduce the risk of hyperplastic changes of the endometrium, patients without a uterus do not require a progestin for this purpose.

### **Usual Dosage Range**

The lowest dose that will control symptoms should be chosen and medication should be discontinued as promptly as possible.

0.5 g to 2 g daily, intravaginally or topically depending on the severity of the condition. Administration should be cyclic (e.g., three weeks on and one week off).

Appropriate diagnostic measures should be taken to rule out malignancy in the event of persistent or recurring abnormal vaginal bleeding.

### **Instructions for Use of Applicator:**

1. Remove cap.
2. Screw nozzle end of applicator onto the tube.
3. Gently squeeze tube to force sufficient cream into the barrel to provide the prescribed dose.
4. Unscrew applicator from tube.
5. Place the applicator into the vaginal opening.
6. To release medication, press plunger downward.

**To Cleanse:** Pull plunger out from barrel. Wash with mild soap and warm water. DO NOT BOIL.

### **Missed Dose**

If a patient misses a dose, it should be taken as soon as possible. If it is close to the patient's next scheduled dose, the missed dose should be skipped, and the patient should continue with her normal schedule. The patient should not take two doses at the same time.

### **Administration**

#### **Vaginal**

Premarin® Vaginal Cream should be prescribed with an appropriate dosage of a progestin for women with intact uteri in order to prevent endometrial hyperplasia/carcinoma. Progestin therapy is not required as part of hormone replacement therapy in women who have had a previous hysterectomy.

### **OVERDOSAGE**

#### **Symptoms of overdose:**

Numerous reports of ingestion of large doses of estrogen products and estrogen-containing oral contraceptives by young children have not revealed acute serious ill effects.

Overdosage with estrogen may cause nausea, breast discomfort, fluid retention, bloating or vaginal bleeding in women.

## **Treatment of overdose**

There is no specific antidote and further treatment if necessary should be symptomatic.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Endogenous estrogens are largely responsible for the development and maintenance of the female reproductive system and secondary sex characteristics. By a direct action, they cause growth and development of the uterus, fallopian tubes, and vagina. With other hormones, such as pituitary hormones and progesterone, they cause enlargement of the breasts through promotion of ductal growth, stromal development, and the accretion of fat. Estrogens are intricately involved with other hormones, especially progesterone, in the processes of the ovulatory menstrual cycle and pregnancy, and affect the release of pituitary gonadotropins. Indirectly, they also contribute to the shaping of the skeleton, maintenance of tone and elasticity through the increase of collagen production in the supportive tissues of the heart, skin and urogenital structures, changes in the epiphyses of the long bones that allow for the pubertal growth spurt and its termination, growth of axillary and pubic hair, and pigmentation of the nipples and genitals. Decline of ovarian estrogenic and progestogenic activity at the end of the menstrual cycle can result in menstruation, although the cessation of progesterone secretion is the most important factor in the mature ovulatory cycle. However, in the preovulatory or anovulatory cycle, estrogen is the primary determinant in the onset of menstruation.

Although circulating estrogens exist in a dynamic equilibrium of metabolic interconversions, estradiol is the principal intracellular human estrogen and is substantially more potent than its metabolites, estrone and estriol at the receptor level. The primary source of estrogen in normally cycling adult women is the ovarian follicle, which secretes 70 to 500 micrograms of estradiol daily, depending on the phase of the menstrual cycle. After menopause, most endogenous estrogen is produced by conversion of androstenedione, secreted by the adrenal cortex, to estrone by peripheral tissues. Thus, estrone and the sulfate conjugated form, estrone sulfate, are the most abundant circulating estrogens in postmenopausal women.

Circulating estrogens modulate pituitary gonadotropins, luteinizing hormone (LH) and follicle stimulating hormone (FSH) through a negative feedback mechanism. Estrogen therapy acts to reduce elevated levels of these hormones seen in postmenopausal women.

Estrogen drug products act by regulating the transcription of a limited number of genes. They may act directly at the cell's surface via non "estrogen receptor" mechanism or directly with the estrogen receptor inside the cell. Estrogens diffuse through cell membranes, distribute themselves throughout the cell, and bind to and activate the nuclear estrogen receptor, a DNA-binding protein which is found in estrogen-responsive tissues. The activated estrogen receptor

binds to specific DNA sequences, or hormone-response elements, which enhance the transcription of adjacent genes and in turn lead to the observed effects. Estrogen receptors have been identified in the wall of blood vessels, in tissues of the reproductive tract, breast, pituitary, hypothalamus, liver, and bone of women.

Conjugated estrogens used in therapy are soluble in water and are well absorbed through the skin, mucous membranes, and the gastrointestinal tract after release from the drug formulation. Some estrogens are excreted in bile; however, they are reabsorbed from the intestine and returned to the liver through the portal venous system. Water-soluble estrogen conjugates are strongly acidic and are ionized in body fluids, which favours excretion through the kidneys since tubular reabsorption is minimal.

Estrogens used in therapy are also well absorbed through the skin and mucous membranes. When applied for a local action, absorption is usually sufficient to cause systemic effects. When conjugated with aryl and alkyl groups for parenteral administration, the rate of absorption of oily preparations is slowed with a prolonged duration of action, such that a single intramuscular injection of estradiol valerate or estradiol cypionate is absorbed over several weeks.

Administered estrogens and their esters are handled within the body essentially the same way as the endogenous hormones.

## **Pharmacodynamics**

Conjugated estrogens used in therapy are soluble in water and are well absorbed through the skin, mucous membranes, and gastrointestinal tract after release from the drug formulation.

### *Effects on vasomotor symptoms associated with estrogen deficiency*

Hot flushes, feelings of intense heat over the upper trunk and face, with flushing of the skin and sweating occur in approximately 80% of women as a result of the decrease in ovarian hormones. These vasomotor symptoms are seen in women whether menopause is surgically induced or spontaneous. However, hot flushes may be more severe in women who undergo surgical menopause. Hot flushes can begin before the cessation of menses.

### *Effects on the Endometrium*

The use of unopposed estrogen therapy has been associated with an increased risk of endometrial hyperplasia, a possible precursor of endometrial adenocarcinoma.<sup>47, 48</sup> The results of clinical studies indicate that the addition of a progestin to an estrogen replacement regimen for more than 10 days per cycle reduces the incidence of endometrial hyperplasia and the attendant risk of adenocarcinoma in women with intact uteri. The addition of a progestin into an estrogen replacement regimen has not been shown to interfere with the efficacy of estrogen replacement therapy for its approved indications.

## **Effect on bleeding patterns**

With a continuous therapy, several bleeding patterns may occur. These may range from absence of bleeding to irregular bleeding. If bleeding occurs, it is frequently light spotting or moderate bleeding.

## **Pharmacokinetics**

### Absorption

Conjugated estrogens are soluble in water and are well absorbed through the skin, mucous membranes, and the gastrointestinal tract after release from the drug formulation.

### Distribution

The distribution of exogenous estrogens is similar to that of endogenous estrogens. Estrogens are widely distributed in the body and are generally found in higher concentration in the sex hormone target organs. Estrogens circulate in the blood largely bound to sex hormone binding globulin (SHBG) and albumin.

### Metabolism

Metabolic conversion of estrogens occurs primarily in the liver (first pass effect), but also at local target tissue sites. Complex metabolic processes result in a dynamic equilibrium of circulating conjugated and unconjugated estrogenic forms which are continually interconverted, especially between estrone and estradiol and between esterified and non-esterified forms.

Estrogen drug products administered by non-oral routes, while not subject to true “first-pass” metabolism, do undergo significant hepatic uptake, metabolism, and enterohepatic recycling. Metabolism and inactivation occur primarily in the liver. Some estrogens are excreted into the bile; however, they are re-absorbed from the intestine and returned to the liver through the portal venous system. Water-soluble estrogen conjugates are strongly acidic and are ionized in body fluids, which favour excretion through the kidneys since tubular re-absorption is minimal.

When given orally, naturally-occurring estrogens and their esters are extensively metabolized (first pass effect) and circulate primarily as estrone sulfate, with smaller amounts of other conjugated and unconjugated estrogenic species. This results in limited oral potency. By contrast, synthetic estrogens, such as ethinyl estradiol and the nonsteroidal estrogens, are degraded very slowly in the liver and other tissues, which results in their high intrinsic potency.

### Excretion

A certain proportion of the estrogen is excreted into the bile, then reabsorbed from the intestine and returned to the liver through the portal venous system. During this enterohepatic recirculation, estrogens are desulfated and resulfated and undergo degradation through

conversion to less active estrogens (estriol and other estrogens), oxidation to nonestrogenic substances (catecholestrogens, which interact with catecholamine metabolism, especially in the central nervous system), and conjugation with glucuronic acids (which are then rapidly excreted in the urine).

### **Special Populations and Conditions**

No pharmacokinetic studies were conducted in special populations, including patients with renal or hepatic impairment.

### **STORAGE AND STABILITY**

Store at 15°C - 30°C.

Keep out of reach of children.

### **SPECIAL HANDLING INSTRUCTIONS**

None required.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

<sup>Pt</sup>PREMARIN® Vaginal Cream is available in tubes of 14 g, each gram containing 0.625 mg of conjugated estrogens CSD. Each tube is accompanied with a calibrated plastic applicator.

Non-Medicinal Ingredients:                   Cetyl Alcohol, Cetyl Esters Wax, Glycerin, Glyceryl Monostearate, Methyl Stearate, Mineral Oil, Phenylethyl Alcohol, Propyl Glycol Monostearate, Sodium Lauryl Sulfate, Water Purified, White Wax.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: Conjugated estrogens, C.S.D.

Chemical name: Not applicable

Molecular formula and molecular mass: Not applicable

Structural formula: Not applicable

Description: Conjugated estrogens C.S.D. contains a mixture of estrogens obtained exclusively from natural sources, occurring as the sodium salts of water-soluble estrogen sulfates blended to represent the average composition of material derived from pregnant mares' urine. It is a mixture of at least the following estrogens: estrone, equilin, 17 $\alpha$ -dihydroequilin, 17 $\alpha$ -estradiol, 17 $\beta$ -dihydroequilin,  $\delta$ 8,9-dehydroestrone, 17 $\beta$ -estradiol, equilenin, 17 $\alpha$ -dihydroequilenin, 17 $\beta$ -dihydroequilenin and as salts of their sulfate esters.

### CLINICAL TRIALS

#### Published Studies

##### *Vasomotor Symptoms and Vaginal Atrophy*

**The Women's Health, Osteoporosis, Progestin, Estrogen (HOPE) Study<sup>44</sup>** was an RCT to evaluate the safety and efficacy of lower doses of CEE and MPA in postmenopausal women. The design included a one year basic study to evaluate the efficacy of lower doses of CEE with and without MPA in relieving vasomotor symptoms (VMS) and vulvar and vaginal atrophy (VVA). A total of 2,673 healthy, postmenopausal women 40 to 65 years of age with an intact uterus (mean age of 53.3 years), including a vasomotor symptom efficacy-evaluable population (n=241 at baseline) participated.

Efficacy measures were frequency and severity of daily hot flushes and Papanicolaou smear with vaginal maturation index (VMI) to assess vaginal atrophy.

There were a total of eight treatment arms consisting of the following:  
CEE 0.625 mg/day; CEE 0.625 mg/MPA 2.5 mg/day; CEE 0.45 mg/day; CEE 0.45 mg/MPA 2.5 mg/day; CEE 0.45 mg/MPA 1.5 mg/day; CEE 0.3 mg/day; CEE 0.3 mg/MPA 1.5 mg/day; or placebo.

Key observations for VMS: All active treatment groups significantly reduced mean number of hot flushes from baseline by week 1 or 2 ( $P<0.01$ ) and all active treatment groups significantly reduced mean number of hot flushes compared with placebo by week 2 or 3 ( $P<0.001$ ).

#### Numbers of hot flushes

- For the placebo group, the mean daily number of hot flushes dropped from approximately 10 at week 1, to approximately 5 at week 12, and continuing at approximately 5 to cycle 13.
- For the 0.625 mg CEE/2.5 mg MPA treatment group, the mean daily number of hot flushes decreased from approximately 10 at week 1, to approximately 1 at week 12, dropping to approximately 0.5 at cycle 13. The difference from placebo was significant ( $P<0.5$ ) beginning from week 2 to the end of cycle 13.

#### Severity of hot flushes

A mild hot flush was rated a 1, a moderate hot flush a 2, and a severe hot flush a 3.

- For the placebo group, the mean daily severity of hot flushes decreased from approximately 2.1 at week 1, to approximately 1.7 at week 12, and continuing at approximately 1.7 to cycle 13.
- For the 0.625 mg CEE/2.5 mg MPA treatment group, the mean daily severity of hot flushes decreased from 2.1 at week 1, to approximately 0.5 at week 12, and dropping to approximately 0.2 at cycle 13. The difference from placebo was significant ( $P<0.5$ ) beginning from week 2 to the end of cycle 13.

Key observations for VVA: All active treatment groups significantly increased the percentage of superficial cells from baseline at cycles 6 and 13 ( $P<0.001$ ) and all active treatment groups significantly increased the percentage of superficial cells compared with placebo at cycles 6 and 13 ( $P<0.001$ ).

## **DETAILED PHARMACOLOGY**

See “Action and Clinical Pharmacology” section under the Health Professional Information Section.

## TOXICOLOGY

Acute toxicity studies have been conducted with conjugated estrogens (Premarin®).

### Acute Toxicity Premarin®

In studies conducted by Wyeth, Premarin (125 mg/kg) was administered orally. The LD<sub>50</sub> value for Premarin® administered orally or intraperitoneally to male and female CD-1 mice and CD rats was greater than 125 mg/kg.

## REFERENCES

### SELECTED BIBLIOGRAPHY

1. Akkad AA, Halligan AW, Abrams K, et al. Differing responses in blood pressure over 24 hours in normotensive women receiving oral or transdermal estrogen replacement therapy. *Obstet Gynecol.* 1997; 89:97-103.
2. Anderson GL, et al. Effects of Estrogen Plus Progestin on Gynecologic Cancers and Associated Diagnostic Procedures. *JAMA.* 2003; 290:1739-1748.
3. Anderson GL, et al. Effects of Conjugated Equine Estrogen in Postmenopausal Women with Hysterectomy. *JAMA.* 2004; 291:1701-1712.
4. Carlström K, et al. Effects of intravaginal oestrogen treatment upon the vaginal absorption of conjugated equine oestrogens. *Maturitas.* 1982; 4:277.
5. Cauley JA, et al. Effects of Estrogen Plus Progestin on Risk of Fracture and Bone Mineral Density. *JAMA.* 2003; 290:1729-1738.
6. Chlebowski RT, Hendrix SL, Langer RD, et al. The Women's Health Initiative Randomized Trial. Influence of Estrogen plus Progestin on Breast Cancer and Mammography in Healthy Postmenopausal Women. *JAMA.* 2003; 289(24):3243-3253.
7. Deutsch S, Ossowski R, and Benjamin I. Comparison between degree of systemic absorption of vaginally and orally administered estrogens at different dose levels in postmenopausal women. *Am. J. Obstet. Gynecol.* 1981; 139(8):967-968.
8. Gelfand MM, and Ferenczy A. A prospective 1-year study of estrogen and progestin in postmenopausal women: Effects on the endometrium. *Obstet Gynecol.* 1989; 74 (3, Part I): 398-402.
9. Grady D, Herrington D, Bittner V, Blumenthal R, Davidson M, Hlatky M, et al. Cardiovascular disease outcomes during 6.8 years of hormone therapy: Heart and Estrogen/progestin Replacement Study follow-up (HERS II). *JAMA.* 2002;288(1):49-57.
10. Gray LA. Treatment of atrophic vaginitis. *J Ky Med Assoc.* 1979; 77:350.
11. Handa VL, Bachus KE, Johnston WW, Robboy SJ, Hammond CB. Vaginal administration of low-dose conjugated estrogens. Systemic absorption and effects on the endometrium. *Obstet & Gynecol.* 1994; 84:215-8.

12. Hassager C, Christiansen C. Blood pressure during oestrogen/progestogen substitution therapy in healthy post-menopausal women. *Maturitas*. 1988; 9(4):315-323.
13. Hintz BL, Kagan AR, Gilbert HA, Rao AR, Chan P, Nussbaum H. Systemic absorption of conjugated estrogenic cream by the irradiated vagina. *Gynecol Oncol*. 1981; 12:75-82.
14. Horwitz RI, and Feinstein AR. Intravaginal estrogen creams and endometrial cancer. No causal association found. *JAMA*. 1979; 241:1266.
15. Hulley S, Furberg C, Barrett-Conner E, et al. Noncardiovascular disease outcomes during 6.8 years of hormone therapy (HERS II). *JAMA*. 2002; 288(1):58-66.
16. Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B, Vittinghoff E. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. *JAMA*. 1998;280(7):605-13.
17. Johnston S., et al. The Detection and Management of Vaginal Atrophy. *J Obstet Gynaecol Can*. 2004; 26(5):503-508.
18. Lacey JV, Mik PJ, Lubin JH, Sherman ME, Troisi R, Hartge P, Schatzkin A, Schainer C. Menopausal hormone replacement therapy and risk of ovarian cancer. *JAMA*. 2002; 288(3): 334-341.
19. Li C, Malone K, Porter P, Weiss N, Tang M, Cushing-Hauzen K, Daling J. Relationship Between Long Durations and Different Regimens of Hormone Therapy and Risk of Breast Cancer. *JAMA*. 2003; 289(24):3254-3263.
20. Lind T, Cameron EC, Hunter WM, et al. A prospective, controlled trial of six forms of hormone replacement therapy given to postmenopausal women. *Br J Obstet Gynaecol*. 1979; 86(3):1-29.
21. Mack TM, et al. Estrogens and Endometrial Cancer in a Retirement Community. *N Eng J Med*. 1976; 294:1262.
22. Mandel FP, Geola FL, Meldrum DR, Lu JHK, Eggena P, Sambhi MP, Hershman JM, et al. Biological effects of various doses of vaginally administered conjugated equine estrogens in post-menopausal women. *J Clin Endocrinol Metab*. 1983; 57:133-9.

23. Manonai J, et al. The Effect of Estradiol Vaginal Tablet and Conjugated Estrogen Cream on Urogenital Symptoms in Postmenopausal Women. *J Obstet Gynaecol Res.* 2001; 27(5): 255-260.
24. Manson JE, et al. Estrogen plus Progestin and the Risk of Coronary Heart Disease. *N Engl J Med.* 2003;349:523-34.
25. Martin PL, Yen SCC, Burnier AM, Hermann H. Systemic absorption and sustained effects of vaginal estrogen creams. *JAMA.* 1979; 242:2699-2700.
26. Million Women Study Collaborators. Breast cancer and hormone-replacement therapy in the Million Women Study. *Lancet.* 2003; 362:419-427.
27. Nathan L. Vulvovaginal disorders in the elderly woman. *Clin Obstet Gynecol.* 1996; 9:933-45.
28. Pickar JH, Yeh I-T, Wheeler JE, Cunnane MF, Speroff L. Endometrial effects of lower doses of conjugated equine estrogens and medroxyprogesterone acetate. *Fertil Steril.* 2001; 76:25-31
29. Raymundo N, et al. Treatment of atrophic vaginitis with topical conjugated equine estrogens in postmenopausal Asian women. *CLIMACTERIC.* 2004; 7: 312-318
30. Rigg LE, Hermann H, and Yen SSC. Absorption of estrogens from vaginal creams. *N Eng J Med.* 1978; 298:195.
31. Rodriguez C, Patel AV, Calle EE, Jacob EJ, Thun MJ. Estrogen replacement therapy and ovarian cancer mortality in a large prospective study of US women. *JAMA.* 2001; 285(11):1460-1465.
32. Shumaker SA, et al. Conjugated Equine Estrogens and Incidence of Probable Dementia and Mild Cognitive Impairment in Postmenopausal Women. *JAMA.* 2004; 291:2947-2958.
33. Shumaker SA, Legault C, Rapp SR, et al. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women – The Women’s Health Initiative Memory Study: a randomized controlled trial. *JAMA.* 2003; 289(20):2651-62.
34. Sitruk-Ware R. Estrogen therapy during menopause. Practical treatment recommendations. *Drugs.* 1990; 39:203-17.
35. Smith DC, et al. Association of Exogenous Estrogen and Endometrial Carcinoma. *N Eng J Med.* 1975; 293:1164-1167.
36. Stewart D. Effect of Premarin Vaginal Cream on latex condoms (US). Wyeth-Ayerst GTR-22769, 1993.

37. Stewart D. Effect of Premarin Vaginal Cream on latex condoms (UK). Wyeth-Ayerst GTR-22770, 1993.
38. The Writing Group for the PEPI Trial. Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. *JAMA*. 1995; 273(3):199-208.
39. The Writing Group for the PEPI Trial. Effects of hormone replacement therapy on endometrial histology in postmenopausal women: The postmenopausal estrogen/progestin interventions (PEPI) trial. *JAMA*. 1996; 275 (5):370-375.
40. Woodruff JD, Pickar JH. Incidence of endometrial hyperplasia in postmenopausal women taking conjugated estrogens (Premarin) with MPA or conjugated estrogens alone. *Am J Obstet Gynecol*. 1994; 170 (5):1213-1223.
41. Wren BG, Routledge AD. The effect of type and dose of oestrogen on the blood pressure of post-menopausal women. *Maturitas*. 1983; 5:135-142.
42. Writing Group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestins in healthy postmenopausal women. *JAMA*. 2002; 288(3):321-333.
43. Ziel HK, Finkle WD. Increased Risk of Endometrial Carcinoma Among Users of Conjugated Estrogens. *N Eng J Med*. 1975; 293:1167.
44. Utian WH, Shoupe D, Bachmann G, Pinkerton JV, and Pickar JH. Relief of vasomotor symptoms and vaginal atrophy with lower doses of conjugated equine estrogens and medroxyprogesterone acetate. *Fertil Steril*. 2001; 75:1065-79.
45. Johnston Jr CC, Melton III LJ, Lindsay R, Eddy DM. Clinical indications for bone mass measurements. *J Bone Miner Res*. 1989;4(suppl 2):1-28.
46. Lafferty FW, Helmuth DO. Post-menopausal estrogen replacement: The prevention of osteoporosis and system effects. *Maturitas*. 1985;7:147-59.
47. Morrow CP, Townsend DE. Cancer of the uterine corpus. In: *Synopsis of gynecologic oncology*. New York: John Wiley and Sons, 2nd ed.; 1981. p. 133-185.
48. Kurman RJ, Kaminsk PF, Norris HJ. The behavior of endometrial hyperplasia: A long-term study of "untreated" hyperplasia in 170 patients. *Cancer*. 1985;56:403-12.

## PART III: CONSUMER INFORMATION

### Premarin® Vaginal Cream (Conjugated Estrogens CSD 0.625 mg/g)

#### IMPORTANT: PLEASE READ

This leaflet is part III of a three-part "Product Monograph" published when Premarin® Vaginal Cream was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Premarin® Vaginal Cream. Contact your doctor or pharmacist if you have any questions about the drug.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

- To treat vulvar and vaginal atrophy and kraurosis vulvae (itching, burning, dryness in or around the vagina, difficulty or burning on urination) associated with menopause.
- To treat dyspareunia (pain during intercourse) associated with menopause.

Premarin® Vaginal Cream should not be used by women with intact uteri unless it is prescribed in association with a progestin.

Premarin® Vaginal Cream should be used only under the supervision of a doctor, with regular follow-up at least once a year to identify side effects associated with its use.

Your first follow-up visit should be within 3 to 6 months of starting treatment. Your visit may include a blood pressure check, a breast exam, a Pap smear and pelvic exam. You should have a mammogram before starting treatment and at regular intervals as recommended by your doctor. Your doctor may recommend some blood tests.

You should carefully discuss the risks and benefits of hormone replacement therapy (HRT) with your doctor. You should regularly talk with your doctor about whether you still need treatment with HRT.

##### What it does:

When using Premarin® Vaginal Cream women are using a hormone, estrogen (i.e. conjugated estrogens CSD 0.625 mg/g). Premarin® Vaginal Cream replaces estrogen specifically in and around the vagina which naturally decreases at menopause.

Estrogens are female hormones that are produced by a woman's ovaries and are necessary for normal sexual development and the regulation of menstrual periods during the childbearing years.

When a woman is between the ages of 45 and 55, the ovaries normally stop making estrogens. This leads to a drop in body estrogen levels and marks the beginning of menopause (the end of monthly menstrual periods). A sudden drop in estrogen levels also occurs if both ovaries are removed during an operation before natural menopause takes place. This is referred to as surgical menopause.

When the estrogen levels begin dropping, some women develop very uncomfortable symptoms, such as feelings of warmth in the face, neck, and chest, or sudden intense episodes of heat and sweating ("hot flashes") as well as vaginal symptoms. In some women the symptoms are mild; in others they can be severe. These symptoms may last a few months or longer.

##### When it should not be used:

Before using Premarin® Vaginal Cream be sure to tell your doctor if you have any of the following medical problems, as Premarin® Vaginal Cream should not be used under these conditions:

- **Known, suspected, or past history of breast cancer.**
- **Known or suspected hormone-dependent cancer.** Estrogens may increase the chances of getting certain types of cancers, including cancer of the breast or uterus. If you have or had cancer, talk with your healthcare provider about whether you should take Premarin® Vaginal Cream.
- **Unexpected or unusual vaginal bleeding**
- **Have (or have had) blood clot disorders, including blood clots in the legs or lungs or thrombophlebitis (inflammation of the veins).**
- **Serious liver disease**
- **Active or past history of heart disease, heart attacks or stroke.**
- **If you are allergic to Premarin® Vaginal Cream or any of its ingredients, or have had any unusual reactions to its ingredients (see What the medicinal ingredients are and What the nonmedicinal ingredients are).**
- **If you are pregnant or suspect you may be pregnant.** Since pregnancy may be possible early in pre-menopause while you are still having spontaneous periods, the use of non-hormonal birth control should be discussed with your physician at this time. If you accidentally take estrogen

during pregnancy, there is a small risk of your unborn child having birth defects.

- If you have partially or completely lost vision due to blood vessel disease of the eye.
- If you have overgrowth of the lining of the uterus.

**What the medicinal ingredients are:**

Conjugated equine estrogens

**What the nonmedicinal ingredients are:**

The cream contains the following inactive ingredients: Cetyl Alcohol, Cetyl Esters Wax, Glycerin, Glyceryl Monostearate, Methyl Stearate, Mineral Oil, Phenylethyl Alcohol, Propyl Glycol Monostearate, Sodium Lauryl Sulfate, Water Purified, White Wax.

**What dosage forms it comes in:**

Premarin® Vaginal Cream is available in tubes of 14 g, each gram containing 0.625 mg of conjugated estrogens CSD. Each tube is accompanied with a calibrated plastic applicator.

You should follow the dosage regimen prescribed by your healthcare provider.

**Instructions for Use of Applicator:**

1. Remove cap.
2. Screw nozzle end of applicator onto the tube.
3. Gently squeeze tube to force sufficient cream into the barrel to provide the prescribed dose.
4. Unscrew applicator from tube.
5. Place the applicator into the vaginal opening.
6. To release medication, press plunger downward.

TO CLEANSE: Pull plunger out from barrel. Wash with mild soap and warm water.

DO NOT BOIL.

**WARNINGS AND PRECAUTIONS**

**Serious Warnings and Precautions**

The Women’s Health Initiative (WHI) is a large clinical study that assessed the benefits and risks of oral combined *estrogen plus progestin* therapy and oral *estrogen-alone* therapy compared with placebo (a pill with no active ingredients) in postmenopausal women.

The WHI trial indicated an increased risk of myocardial infarction (heart attack), stroke, breast cancer, pulmonary emboli (blood clots in the lungs) and deep vein thrombosis (blood clots in the large veins) in postmenopausal women taking oral combined *estrogen plus progestin*.

The WHI trial indicated an increased risk of stroke and deep vein thrombosis in postmenopausal women with prior hysterectomy (surgical removal of the uterus) taking oral *estrogen-alone*.

Therefore, you should highly consider the following:

- There is an increased risk of developing invasive breast cancer, heart attack, stroke and blood clots in both lungs and large veins with the use of estrogen plus progestin therapy.
- There is an increased risk of stroke and blood clots in the large veins with the use of estrogen-alone therapy.
- Estrogens with or without progestins should not be used for the prevention of heart disease or stroke.
- Estrogens with or without progestins should be used at the **lowest effective dose** and for the **shortest period of time** possible. Regular medical follow-up is advised.

**Breast Cancer**

The results of the WHI trial indicated an increased risk of breast cancer in postmenopausal women taking combined *estrogen plus progestin* compared to women taking placebo.

The results of the WHI trial indicated no difference in the risk of breast cancer in post-menopausal women with prior hysterectomy taking *estrogen-alone* compared to women taking placebo.

Estrogens should not be taken by women who have a personal history of breast cancer.

In addition, women with a family history of breast cancer or women with a history of breast lumps, breast biopsies or abnormal mammograms (breast x-rays) should consult with their doctor before starting HRT.

Women should have a mammogram before starting HRT and at regular intervals during treatment as recommended by their doctor.

Regular breast examinations by a doctor and regular breast self-examination are recommended for all women. You should review technique for breast self-examination with your doctor.

### ***Overgrowth of the lining of the uterus and cancer of the uterus***

The use of *estrogen-alone* therapy by post-menopausal women who still have a uterus increases the risk of developing endometrial hyperplasia (overgrowth of the lining of the uterus), which increases the risk of endometrial cancer (cancer of the lining of the uterus).

If you still have your uterus, you should take a progestin medication (another hormone drug) regularly for a certain number of days of each month to reduce the risk of endometrial hyperplasia.

You should discuss progestin therapy and risk factors for endometrial hyperplasia and endometrial carcinoma with your doctor. You should also report any unexpected or unusual vaginal bleeding to your doctor.

If you have had your uterus removed, you are not at risk of developing endometrial hyperplasia or endometrial carcinoma. Progestin therapy is therefore not generally required in women who have had a hysterectomy.

### ***Heart Disease and Stroke***

The results of the WHI trial indicated an increased risk of stroke and coronary heart disease in post-menopausal women taking combined *estrogen plus progestin* compared to women taking placebo.

The results of the WHI trial indicated an increased risk of stroke, but no difference in the risk of coronary heart disease in post-menopausal women with prior hysterectomy taking *estrogen-alone* compared to women taking placebo.

### ***Abnormal Blood Clotting***

The results of the WHI trial indicated an increased risk of blood clots in the lungs and large veins in post-menopausal women taking combined *estrogen plus progestin* compared to women taking placebo.

The results of the WHI trial indicated an increased risk of blood clots in the large veins, but no difference in the risk of blood clots in the lungs in post-menopausal women with prior hysterectomy taking *estrogen-alone* compared to women taking placebo.

The risk of blood clots also increases with age, if you or a family member has had blood clots, if you smoke or if you are severely overweight. The risk of blood clots is also temporarily increased if you are immobilized for long periods of time and following major surgery. You should discuss risk factors for blood clots with your doctor since blood clots can be life-threatening or cause serious disability.

### ***Gallbladder Disease***

The use of estrogens by postmenopausal women has been associated with an increased risk of gallbladder disease requiring surgery.

### ***Dementia***

The Women's Health Initiative Memory Study (WHIMS) was a substudy of the WHI trial and indicated an increased risk of dementia (loss of memory and intellectual function) in post-menopausal women age 65 and over taking oral combined *estrogen plus progestin* compared to women taking placebo.

The WHIMS indicated no difference in the risk of dementia in post-menopausal women age 65 and over with prior hysterectomy taking oral *estrogen-alone* compared to women taking placebo.

### **BEFORE you use Premarin® Vaginal Cream talk to your doctor or pharmacist if you:**

- have a history of allergy or intolerance to any medications or other substances
- have a personal history of breast disease (including breast lumps) and/or breast biopsies, or a family history of breast cancer
- have experienced any unusual or undiagnosed vaginal bleeding
- have a history of uterine fibroids or endometriosis
- have a history of liver disease, jaundice (yellowing of the eyes and/or skin) or itching related to estrogen use or during pregnancy
- have a history of migraine headache
- have a history of high blood pressure
- have a personal or family history of blood clots, or a personal history of heart disease or stroke
- have a history of kidney disease, asthma or epilepsy (seizures)
- have a history of bone disease (this includes certain metabolic conditions or cancers that can affect blood levels of calcium and phosphorus)
- have been diagnosed with diabetes
- have been diagnosed with porphyria (a disease of blood pigment)
- have a history of high cholesterol or high triglycerides
- are pregnant or may be pregnant
- have had a hysterectomy (surgical removal of the uterus)
- smoke

Other existing conditions you should discuss with your health professional include lupus, very low calcium levels, thyroid problems, fluid retention, gallbladder disease, depression, and breastfeeding. If you have upcoming surgery or prolonged bedrest, you should also discuss these.

## INTERACTIONS WITH THIS MEDICATION

Tell your doctor or pharmacist if you are taking any other medications, including prescription medications, over-the-counter medications, vitamins or herbal products (such as St. John's wort). Some medications (such as medications for high blood pressure, diabetes, blood clots, sleeping, anxiety, seizures, pain-relief and tuberculosis) may affect how Premarin® Vaginal Cream works. Premarin® Vaginal Cream may also affect how other medicines work.

## PROPER USE OF THIS MEDICATION

### Usual dose:

You should follow the dosage regimen prescribed by your healthcare provider.

Estrogens should be used at the lowest dose possible for your treatment only as long as needed. You and your healthcare provider should talk regularly (for example every 3 to 6 months) about the dose you are taking and whether you still need treatment with Premarin® Vaginal Cream.

Do not give Premarin® Vaginal Cream to other people, even if they have the same symptoms you have. It may harm them.

### Overdose:

Contact your physician or local Poison Control Center in case of accidental ingestion of high doses of Premarin® Vaginal Cream.

**Overdosage with estrogens may cause nausea and vomiting, breast discomfort, fluid retention, bloating or vaginal bleeding may occur in women. There is no specific antidote and further treatment if necessary should be symptomatic.**

Overdosage may result in a period of amenorrhea (lack of menses) of a variable length and may be followed by irregular menses for several cycles. No cases of overdosage in male patients have been reported.

### Missed Dose:

If you miss a dose, use it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your normal schedule. Do not use 2 doses at the same time.

## SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Possible side effects include:

Breast pain; breakthrough bleeding; spotting; change in menstrual flow; a spontaneous flow of milk from the nipple; painful periods; nausea; bloating; abdominal pain; vomiting; dizziness; headache (including migraine); joint pain; changes in libido; mood disturbances; irritability; hair loss; rash; itching; tender red nodules on the shins and legs; inflammation of the vagina; hives; changes in weight (increase or decrease); increase in blood pressure

## SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / possible side effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Abdominal pain, nausea or vomiting	✓		
Breast lump, unusual discharge		✓	
Shortness of breath, weakness, unusual fatigue, cold sweat, dizziness, sleep disturbance, indigestion, anxiety		✓	
Pain or swelling in the leg			✓
Persistent sad mood		✓	
Sharp pain in the chest, coughing blood or sudden shortness of breath			✓
Sudden partial or complete loss of vision			✓
Sudden severe headache or worsening of headache, vomiting, dizziness, fainting, disturbance of vision or speech or weakness or numbness in an arm or leg			✓
Unexpected vaginal bleeding		✓	
Yellowing of the skin or eyes		✓	

This is not a complete list of side effects. For any unexpected effects while taking Premarin® Vaginal Cream, contact your doctor or pharmacist.

## HOW TO STORE IT

Store Premarin® Vaginal Cream at 15° C to 30° C (room temperature).

Keep out of reach of children.

### **REPORTING SUSPECTED SIDE EFFECTS**

To monitor drug safety, Health Canada collects information on serious and unexpected effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Health Canada by:

toll-free telephone: 866-234-2345

toll-free fax: 866-678-6789

By email: [cadrmp@hc-sc.gc.ca](mailto:cadrmp@hc-sc.gc.ca)

By regular mail:

Canadian Adverse Drug Reaction Monitoring Program  
(CADRMP)

Marketed Health Products Directorate

Health Canada

Tunney's Pasture, Address Locator: 0701C

Ottawa ON K1A 0K9

*NOTE: Before contacting Health Canada, you should contact your physician or pharmacist.*

### **MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals can be found at:

<http://www.wyeth.ca/en>

or by contacting the sponsor, Wyeth Canada, at:

1-800-461-8844

This leaflet was prepared by Wyeth Canada

Last revised: June 01, 2007